Dr. Travis Frantz; Leslie Solis, PA
Texas Health Orthopedic Specialists
2008 L. Don Dodson Drive, Suite 100
Bedford, TX 76021
(817) 283-0967
www.frantzorthopedics.com



Distal Radius Fracture Rehabilitation Protocol (non-operative management)

• Weeks: 0-2

- Goals: avoid fracture displacement, allow early bone healing, decrease swelling/inflammation
- Splint/cast to remain in place
- Non-weightbearing with upper extremity
- Start formal therapy 2-6 weeks from date of surgery, at discretion of provider based upon fracture and type of immobilization required
- Encourage ROM of fingers and shoulder (also elbow if not immobilized in cast/splint)
- Modalities as needed for muscle stimulation, pain control, swelling control

• Weeks 2-6

- Goals: avoid fracture displacement, allow continued bone healing, decrease swelling/inflammation
- Initial cast/splint removed and transitioned into removal Velcro wrist splint or cast at provider's discretion
- Maintain non-weightbearing with extremity; okay for using phone, keyboard, etc. as needed while in splint/cast; nothing more or heavier
- Continue ROM of fingers and shoulder (elbow also if not immobilized in cast/splint)
- Modalities as needed for muscle stimulation, pain control, swelling control

Weeks 6-12

- Goals: improved swelling and pain, increase ROM
- Continue to wear removable Velcro splint when not working on ROM or in therapy
- o Begin gentle active wrist ROM in all planes in week 6, passive to begin in week 10
- Weightbearing/lifting restriction of 5 lbs.; okay to increase to 10 lbs. in week 10
- Emphasize importance of home exercise/ROM program
- Modalities as needed for muscle stimulation, pain control, swelling control

Weeks 12+

- Goals: increase ROM, develop strength, return to work, sport, and/or pre-injury functional level with minimal to no restrictions
- Begin to slowly wean from removable Velcro splint, and eventually discontinue when comfortable doing so
- Continue improving wrist ROM in all planes
- Continue advancing weightbearing/lifting restriction as tolerated
- Work on improving functional/ADL tasks as needed
- Functional/work-specific/sport strengthening and training, if applicable
- Customized HEP to continue once PT/OT is completed
 - Full functional return takes 6 months to 1 year

